



Data Mining of ESEL and JET Probe Data

Hoffmann, Mathias; Naulin, Volker; Nielsen, Anders Henry; Madsen, Jens; Juul Rasmussen, Jens

Publication date:
2009

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hoffmann, M. (Author), Naulin, V. (Author), Nielsen, A. H. (Author), Madsen, J. (Author), & Juul Rasmussen, J. (Author). (2009). Data Mining of ESEL and JET Probe Data. Sound/Visual production (digital)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Data Mining of ESEL & JET probe data

Mathias Hoffmann, Volker Naulin, Anders Nielsen, Jens
Madsen, Jens Juul Rasmussen

Association EURATOM - Risø DTU, Roskilde, Denmark

- Fusion devices: Turbulence
- Internal plasma dynamic: Statistical properties
- Link: plasma turbulence and K41 model ?
- Universal properties of edge magnetized plasmas: Fluctuations are self-similar in space & time

- Fusion devices: Turbulence
- Internal plasma dynamic: Statistical properties
- Link: plasma turbulence and K41 model ?
- Universal properties of edge magnetized plasmas: Fluctuations are self-similar in space & time due to intermittent structures
 - Memory effects
 - Large-scale correlation (anomalous diffusion)
- coherent structures (vortices, zonal flows, streamers, blobs)

- Introduction: SOC & Self-Similarity
- Methods to determine Hurst Coefficient
- JET: Hurst vs. ρ
- ESEL: Hurst vs. ρ

Introduction: Self-Organized Criticality (SOC)

- Phenomena like: self-organization of NLDS + near-critical
length scale : fractal geometry
time scale : power spectra, behavior $1/f$
- SOC systems = scale invariant in space and time
- Experiments: intermittency and self-similar fluctuations
- Memory-effects and long-range correlations (persistence)
- Hurst parameter: suitable measure of persistence in time series

Statistical sense : $x(bt) \stackrel{d}{=} b^H x(t) \quad \forall b, t > 0$

Statistical sense : $x(bt) \stackrel{""}{=} b^H x(t) \quad \forall b, t > 0$

Hurst exponent :

$H = 0.0$ White noise

$H < 0.5$ anti-correlation (restriction of the noise process)

$H = 0.5$ Brownian noise

$H > 0.5$ persistence, fractal Brownian process

$H = 1.0$ exactly self-similar process

stationary process fractional Gaussian noise (fGn)
autocorrelation coefficient

$$\rho_n(X_t, X_{t+n}) = 0.5 \cdot (|n+1|^{2H} - 2|n|^{2H} + |n-1|^{2H})$$

nonstationary process fractal Brownian motion (fBm)
cumulative sum of a fGn process $Y_t = \sum_{t_0}^t X_i$

The processes do not share the same correlation structure!

- Rescaled range (R/S) method, time lag $> 5 - 10$ local autocorrelation time
- Fourier spectra (just monofractal)
- Structurfunctions, time lag > 1 local autocorrelation time
- Scaled window variance method

Power Law Scaling, Wiener Khinchine Theorem link H with β

$$\lim_{\omega \rightarrow \infty} S(\omega) \sim \omega^{-\beta}$$

stationary process $1 < \beta = 2H + 1 < 3$

nonstationary process $-1 < \beta = 2H - 1 < 1$ (time average PSD)

Note: $H = H(q = 2)$ thus only monofractal analysis

Scaled Windows Variance Method (SWVM) for fBm

$$V(m) = \frac{m}{N} \sum^{N/m} \sigma_i \sim m^H$$

Note: $H = H(q = 1)$ thus only monofractal analysis

Example

K41 Theory each point of the \vec{v} -field: $\Delta v_L \sim L^{1/3}$
energy spectrum, no viscosity, $E_k \sim k^{-5/3}$

Example

K41 Theory each point of the \vec{v} -field: $\Delta v_L \sim L^{1/3}$
energy spectrum, no viscosity, $E_k \sim k^{-5/3}$

Structure Functions (SF) for fBm

$$S_q(\Delta t) = \langle |x(t + \Delta t) - x(t)|^q \rangle \sim \Delta t^{\xi(q)}$$

Example

K41 Theory each point of the \vec{v} -field: $\Delta v_L \sim L^{1/3}$
energy spectrum, no viscosity, $E_k \sim k^{-5/3}$

Structure Functions (SF) for fBm

$$S_q(\Delta t) = \langle |x(t + \Delta t) - x(t)|^q \rangle \sim \Delta t^{\xi(q)}$$

Measure of self-similarity Hurst $H = E + 1 - D = \xi(q)/q$

Example

K41 Theory each point of the \vec{v} -field: $\Delta v_L \sim L^{1/3}$
energy spectrum, no viscosity, $E_k \sim k^{-5/3}$

Structure Functions (SF) for fBm

$$S_q(\Delta t) = \langle |x(t + \Delta t) - x(t)|^q \rangle \sim \Delta t^{\xi(q)}$$

Measure of self-similarity Hurst $H = E + 1 - D = \xi(q)/q$

Integrated Structure Functions (intSF) for fGn

$$W_t = \sum_i^t X_i$$

$$\text{int}S_q(\Delta t) = \langle |W(t + \Delta t) - W(t)|^q \rangle \sim \Delta t^{\xi(q)}$$

Measure of self-similarity Hurst $H = E + 1 - D = \xi(q)/q$

Monofractal

$$\xi(q) = Hq$$

Multifractal

$$\xi(q) = H(q)q$$

Measure of self-similarity Hurst $H = E + 1 - D = \xi(q)/q$

Monofractal

$$\xi(q) = Hq$$

Multifractal

$$\xi(q) = H(q)q$$


Intermittent

She and Leveque (SL) predicting scaling model

$$\xi(q) = \frac{q}{9} + 2 \left(1 - \left(\frac{2}{3} \right)^{q/3} \right)$$

- Example: ESEL, structure functions, integrated structure functions (`gnu/den_sat3x3_temperature_data.gnu`)
- Hurst vs. radial direction
(`gnu/den_sat3x3_temperature_all_H.gnu`)
- $\xi(q)$ vs. q
(`gnu/den_sat3x3_temperature_anstiege.gnu`)

- Langmuir-probe measurement: ion saturation current $j_{\text{sat}} \propto n \cdot \sqrt{T_e}$.
- density fluctuation behavior is dominant
- Windows size $\Delta\rho = [0.25, 0.5, 1.0]$



Electrostatic interchange model


$$\frac{D}{Dt}n = \Lambda_n n - n\mathcal{C}(\phi) + \mathcal{C}(nT_e)$$

$$\frac{D}{Dt}T_e = \Lambda_{T_e} T_e - \frac{2}{3}T_e\mathcal{C}(\phi) + \frac{7}{3}T_e\mathcal{C}(T_e) + \frac{2}{3}\frac{T_e^2}{n}\mathcal{C}(n)$$



$$\frac{D}{Dt}\Omega = \Lambda_\Omega \Omega + \mathcal{C}(nT_e)$$

with

$$\mathcal{C}(\cdot) = -\frac{\rho_{s,0}}{R_0} \frac{\partial}{\partial y} \cdot, \quad \Lambda_\cdot = D_\perp \cdot \nabla_\perp^2 \cdot - \frac{\cdot}{\tau_\parallel}, \quad \frac{D}{Dt} = \frac{\partial}{\partial t} + v_E \cdot \nabla \phi$$



ESEL - Probe Movement



- Confined plasma: Persistence dominate the dynamics, transport processes
- SOL: Random processes dominate the dynamics $H \approx 0.5$
- Wall Shadow: High Hurst coefficient due to low statistics

Experiment : $H \approx 0.8$

Simulation : $H \approx 0.7$

- Hurst coefficient in dependence of the shear flow
- Which parameter Multifractality in the simulation?
- Persistence of potential fluctuations are different from j_{sat} ?
- Find jet shots with ESS to study turbulent models ...
- Is a 3D code more suitable?

Thanks for your attention!